




Envu Farm Hygiene Solutions

Products	Active	Group(s)	Key Features	Pest	Outdoor	Indoor	Dilution Rates	Coverage per Container	Re-entry
 ANNIHILATOR PolyZone	Annihilator PolyZone (Deltamethrin 50 g/L) Unit - 473 mL bottle (12 x 473 mL case)	3	Contains deltamethrin, a type II pyrethroid in a unique proprietary polymer suspension. Designed to resist weathering outdoors and lasts up to 90 days. Holds up well indoors in dirty, greasy environments.	Darkling beetles (lesser mealworm), flies (including house flies, blue bottle flies, cluster flies and flesh flies), mosquitoes and ticks	Broadcast, spot and crack-and-crevice treatment on exterior structural surfaces of structure as well as perimeter 1 m out from and 1 m up structure.	Spot treatment and crack and crevice treatment.	6 mL/L or 2 mL/L (maintenance rate) Apply 7.5 L of diluted material to 100 m ² area	1890 m ² (20,234 ft ²) or 5910 m ² (63,614 ft ²)	DO NOT apply when people, pets or livestock are present. DO NOT allow people, pets or livestock to enter treated areas until sprays have dried.
 CREDO SC	Credo SC Insecticide (526 g/L imidacloprid) Unit - 500 mL bottle (8 x 500 mL case)	4	Enhanced efficacy and residual control.	Darkling beetles (larva and adults)	Indoor only	Broadcast, band, spot, crack-and-crevice (floors, walls, support beams)	Dilute 90 mL in 2-7.5 L of water. Apply the diluted mixture to 100 m ² area	555 m ²	DO NOT apply when birds are present. Apply between flocks following de-caking/sanitation. Allow treated surfaces to dry before restocking/reintroducing birds into the facility. DO NOT enter or allow worker entry into treated areas until sprays have dried.
 MAXFORCE Fly Spot Bait	Imidacloprid 10%, Muscalure .1% 56.7 g pouch (50 x 56.7 g case)	4	Combines powerful attractants with imidacloprid (Group 4 insecticide). Applied as a spot treatment only. Flies in close proximity are drawn to bait. Starts to work in 60 seconds once flies land on the bait spots.	House flies	Limited outdoor use on the exterior surfaces of buildings (see label for details). May also be applied to the interior and exterior surfaces of dumpsters and other waste containers.	Agricultural livestock production facility structures such as poultry houses, turkey and chicken houses, cattle, sheep and goat facilities, feedlots, dairy barns, swine confinement buildings, dog kennels, horse stables, animal confinement areas, animal holding areas, animal quarters, animal rearing facilities, livestock barns, hobby and recreational farms and areas of animal facilities including storage rooms, dumpsters and trash collection areas, where house flies tend to aggregate or be a nuisance.	Dilute 1 x 56 g pack per 450 mL of water. Apply 2 or 4 L of diluted mixture to treat 100 m ² of fly resting areas	450 mL of diluted mixture covers 45 m ² of fly resting areas	DO NOT apply when people or pets or livestock are present. DO NOT allow people or pets to re-enter treated areas until sprays have dried.

