Document Date: 8/25/2025
Expiration Date: 12/31/2029

AMINOCYCLOPYRACHLOR	GROUP	4	HERBICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF: TX

Method® 240SL Herbicide

EPA Reg. No. 101563-177

For control of broadleaf and brush species, including invasive species, in non-crop areas.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

METHOD® 240SL HERBICIDE is a soluble liquid that is mixed in water and applied as a spray. METHOD® 240SL HERBICIDE may be applied by aerial or ground equipment for control of broadleaf weeds and woody species, including many terrestrial and riparian invasive and noxious weeds. METHOD® 240SL HERBICIDE is registered for weed and brush control on private, public, and military lands as follows: non-crop areas such as airports, highways/roadsides, railroad, pipeline and utility rights-of-way, sewage disposal areas, industrial areas, such as electrical substations, rail yards or other industrial rock areas, farmyards, fuel storage areas, fence rows (including fences that border grazed areas), non-irrigation ditch banks, barrier strips, lumberyards, pumping stations and tank farms, restoration areas, natural areas, wildlife management areas, wildlife openings, and wildlife habitats. METHOD® 240SL HERBICIDE may be used for the release or restoration of native perennial grasses and in established industrial turf grasses.

METHOD® 240SL HERBICIDE provides preemergence and/or postemergence control of the broadleaf weeds, vines, and brush species listed in the WEEDS CONTROLLED section of the label. For perennial species on the label, a postemergence application should be used. For best postemergence performance, a methylated seed oil (MSO) adjuvant should be included in the spray solution. Excessive wetting of the target plant is not necessary but good spray coverage of the target plant is needed for best results. Weeds hardened off by cold weather or drought stress may not be controlled.

IMPORTANT PRECAUTIONS

Injury or loss of desirable trees or vegetation may result if Method 240SL herbicide is applied on or near desirable trees or vegetation, on areas where their roots extend, in locations where the treated soil may

Environmental Science US, LLC.
5000 CentreGreen Way,
Suite 400
Cary, NC 27513

be washed or moved into the contact with their roots, or where roots of treated plants and roots of nearby desirable trees extend into the same root zone. Certain species may be sensitive to low levels of Method 240SL including but not limited to conifers (such as Douglas fir, Norway spruce, and ponderosa and white pine), deciduous trees (such as aspen, cottonwood, magnolia, poplar, maple species, and willow species), and ornamental shrubs (such as arborvitae, burning bush, crape myrtle, and forsythia).

IMPORTANT RESTRICTIONS

Do not apply Method 240SL within the root zone of desirable trees and/or shrubs unless injury or loss can be tolerated. Root zones of desirable trees/shrubs may extend beyond the tree canopy.
Do not use plant material treated with Method 240SL for mulch or compost.
Do not apply more than 18 fluid ounces of Method 240SL per acre per year on a broadcast-acre basis.

ADDITIONAL INFORMATION

Refer to the Method 240SL label for additional information on use directions, application methodology, adjuvants, restrictions, and precautions.

USE RATES

Method 240SL should be applied at the recommended rate ranges for weeds listed below. Higher rates should be used for larger, established plants or when the density of the target species is high. Do not exceed 18 fl oz/A of Method 240SL in a 12-month period.

Species	Recommended Rate (fl oz/A)	Amount of Method 240SL per 100 gallons of spray solution (fluid ounces)
African Rue (*Peganum harmala*) Common Tansy (*Tanacetum vulgare*) Wild Parsnip (*Pastinaca sativa*) Yellow Toadflax (*Linaria vulgaris*)	8-18	32-72
Big Sagebrush (*Artemisia tridentata*) Buckbrush (*Ceoanthus cuneatus*) Bush Honeysuckle (*Lonicera* spp.) Douglas Fir (*Pseudotsuga menziesii*) Giant Hogweed (*Heracleum mantegazzianum*) Grand Fir (*Abies grandis)* Hemp Dogbane (*Apocynum cannabinum*) Lodgepole Pine (*Pinus contorta*) Mimosa (*Albizia julibrissin*) White Pine (*Pinus strobus*)	12-18	48-72

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-424-9300.

For PRODUCT USE Information Call 1-800-331-2867).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-800-331-2867

Method 240SL is a registered trademark of Environmental Science US LLC.

Environmental Science US, LLC.
5000 CentreGreen Way,
Suite 400
Cary, NC 27513