



TRIPLE CROWN[®]

T&O

Insecticide

For use on lawns and landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

envu[™]

Active Ingredients:

	By Wt.
Zeta-Cypermethrin*	2.70%
Bifenthrin**	7.87%
Imidacloprid	13.83%
Other Ingredients***:	75.60%
	100.0%

*Cis/trans isomer ratio: Max 75% (±) cis and Min. 25% (±) trans

**Cis isomers 97% minimum; trans isomers 3% maximum.

***Contains Petroleum Distillates

TRIPLE CROWN[®] T&O Insecticide contains 2.25 pounds active ingredients per gallon.

EPA Reg. No. 101563-352

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See other panels for additional precautionary information.

SKU: D00001605
22062313B 250407AV1
Net Contents: 1 Gallon

PRODUCED FOR
Environmental Science U.S., LLC
5000 CentreGreen Way, Suite 400
Cary, NC 27513

FIRST AID

IF SWALLOWED

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give **any** liquid to the person.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

This product contains a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Call Environmental Science U.S., LLC Emergency Response Telephone No. 1-800-424-9300 for Emergency Assistance.

For information regarding the use of this product call 1-800-331-2867.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Warning

May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear a long-sleeved shirt and long pants, socks, shoes and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

User Safety Recommendations

Wash thoroughly with soap and water after handling. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help to avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops, **plants** or weeds. Do not apply this product or allow it to drift to blooming crops, **plants or weeds** if bees are foraging in the treatment area.

The chemical imidacloprid demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- o Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- o Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- o Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- o Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens, parks, lawns, or grounds.

Also for use as a broadcast or spot treatment to control pests outdoors, including areas surrounding residential, institutional, public, commercial and industrial buildings.



Restrictions

Do not apply TRIPLE CROWN® T&O insecticide while bees are foraging. Do not apply TRIPLE CROWN T&O insecticide to plants that are flowering. Only apply after all flower petals have fallen.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply by air.

Do not apply this product through any type of irrigation system.

Do not allow people or pets in treated area until spray has dried.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

Do not make applications during rain.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Not for use on plants being grown for sale, or other commercial use, or for commercial seed production, or for research purposes.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

Do not apply more than 0.88 lb ai/A (50 fl oz/A) or 0.50 lb ai imidacloprid/A per year.

Resistance Management

Some insects are known to develop resistance to products with the same chemical class used repeatedly for control. TRIPLE CROWN T&O insecticide contains Group 3 and Group 4A insecticides. Although pest resistance cannot be predicted, a general rule to reduce the onset of resistance in pest species to TRIPLE CROWN T&O insecticide is not to consecutively and repeatedly apply Group 3 and/or Group 4A insecticides during a growing season for control of a particular pest target. Consult your local or state agricultural authorities or your Environmental Science U.S., LLC representative for more specific details on insect resistance management strategies. If resistance does occur use another chemistry class that is registered for the pest and crop.

APPLICATION INSTRUCTIONS

Apply TRIPLE CROWN T&O insecticide as a broadcast or spot treatment.

Use application volumes of up to 10 gallons of final dilution per 1,000 sq ft when treating dense grass or foliage for uniform coverage.

For low volume applications (< 2 gallons/1,000 sq ft), irrigate the treated area with at least 0.25 inches of water immediately following application to ensure efficacy against sub-surface pests (e.g. mole crickets).

TURF APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, TRIPLE CROWN T&O insecticide may be applied at up to 0.80 fl oz per 1,000 sq ft (35 fl oz/A) to control each of the pests listed in this Table. The higher application rates within the rate range should be used when maximum residual control is desired or heavy pest populations exist. Do not apply more than 0.88 lb ai/A (50 fl oz/A) of TRIPLE CROWN T&O insecticide **or 0.50 lb ai of imidacloprid/A** per year.

Pest	Application Rate	
	fl oz/1,000 sq ft	fl oz/A
Armyworms ¹ Cutworms ¹ Sod Webworm ¹	0.23 - 0.34	10 - 15
Annual Bluegrass Weevil (Hyperodes) (Adult) ² Banks Grass Mite ⁵ Billbugs (Adult) ³ Black Turfgrass Ataenius (Adult) ⁴ Centipedes Crickets Earwigs Fleas (Adult) Grasshoppers Leafhoppers Mealybugs Millipedes Mites ⁵ Pillbugs Sowbugs	0.23 - 0.46	10 - 20

(continued)

Pest	Application Rate	
	fl oz/1,000 sq ft	fl oz/A
Ants Black Vine Weevil (Adult) Chinch Bugs ⁶ Crane Flies ⁷ European Chafer Fleas (Larvae) ⁸ Frit Fly Green June Beetle May or June Beetle Imported Fire Ants ⁹ Japanese Beetle (Adult) Masked Chafer (Northern or Southern) Mole Cricket (Adult) ¹⁰ Mole Cricket (Nymph) ¹¹ Stink Bugs Ticks ¹²	0.46 - 0.80	20 - 35

Restrictions:

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of TRIPLE CROWN T&O insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Comments

¹**Armyworms, Cutworms and Sod Webworms:** To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application. If the grass area is being maintained at a mowing height of greater than 1 inch, then higher listed application rates (Up to 0.80 fl oz per 1,000 sq ft) may be required during periods of high pest pressure.

²**Annual Bluegrass Weevil (*Hyperodes*) adults:** Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when *Forsythia* is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

³**Billbug adults:** Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

- 4 Black Turfgrass Ataenius adults:** Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).
- 5 Mites:** To ensure optimal control of eriophyid mites, apply in combination with the labeled application rate of a surfactant. A second application, five to seven days after the first, may be necessary to achieve acceptable control.
- 6 Chinch Bugs:** Chinch Bugs infest the base of grass plants and are often found in the thatch layer. Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located. Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained. Chinch Bugs can be one of the most difficult pests to control in grasses and the higher application rates (Up to 0.8 fl oz per 1,000 sq ft) may be required to control populations that contain both nymphs and adults during the middle of the summer.
- 7 Crane Flies:** Treatments can be made to control early to mid-season larvae (approximately August - February) as they feed on plant crowns. Treatments made to late-season larvae (approximately March, April) may only provide suppression.
- 8 Flea larvae:** Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Use a higher volume application of final dilution when treating these areas to ensure penetration of the insecticide into the soil. Note: if the lawn area is being treated with TRIPLE CROWN T&O insecticide at 0.25 fl oz per 1,000 sq ft for adult flea control, then the larval application rate may be achieved by increasing the application volume two- to four-fold.
- 9 Imported Fire Ants:** Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application or use a high volume application of final dilution. Apply broadcast treatments at 0.46 to 0.80 fl oz per 1,000 sq ft. Use enough finished volume to penetrate thatch or sod. Treat mounds by applying 0.80 fl oz TRIPLE CROWN T&O insecticide per mound in 1 to 2 gallons water by sprinkling the mound until it is wet and treat 3 feet out around the mound. Use the higher volume for mounds larger than 12". Treat mounds with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. For best results, apply in cool weather (65 - 80°F) or in early morning or late evening hours.
- 10 Mole Cricket adults:** Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be

maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

¹¹Mole Cricket nymphs: Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging, nymphs later in the year may require both higher listed application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

¹²Ticks (Including ticks that may transmit Lyme Disease and Rocky Mountain Spotted fever): Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed activity. Limit repeat application to no more than once per seven days.

Deer ticks (*Ixodes* sp.) have a complicated life cycle that ranges over a two year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs and adults.

ORNAMENTAL APPLICATION RATES

Foliar Applications:

Apply at the specified application rate as a full coverage foliar spray. Repeat treatment as needed to achieve control. Use higher labeled application rates as pest pressure and amount of foliage increases.

Broadcast Applications:

For foliage plants and ground covers, mix the required amount of TRIPLE CROWN T&O insecticide in sufficient water to uniformly and accurately cover the treatment area. Do not use less than 2 gallons of water per 1,000 sq ft. For optimum control of soil-borne pests, irrigate thoroughly after treatment to help incorporate TRIPLE CROWN T&O insecticide into the upper layers of soil.

Application to Trees, Shrubs, Evergreens, Foliage plants, and Ground covers**For best results, treatment should be started prior to heavy infestation or pest pressure.**

Pests			Rate fl oz per 100 gallons
Ants Beet Armyworm Black vine weevil adult Broad mites Budworms Scale crawlers Citrus thrips Clover mites ⁵ Diaprepes (adults)	European red mites Flea beetles Fungus gnats (adults) Grasshoppers Imported Fire Ants ⁹ Leafrollers Mites ⁵ Mosquitoes Orchid weevil	Pine needle scales (crawlers) Plant bugs San Jose scale (crawlers) Spider mites Thrips Tip moths Twig borers Wasps	7.5
Bagworms Cutworms	Fall webworms Gypsy moth caterpillars	Leaf feeding caterpillars Tent caterpillars	4 - 7.5
Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles)	Leafhoppers (including glassy-winged sharpshooter) Mealybugs Psyllids	Sawfly larvae Thrips (suppression) Treehoppers Whiteflies	2.5 - 7.5
Restrictions: Target treatment to areas where pest are active. Use sufficient water volume to thoroughly wet plant surfaces. Repeat applications may be made if there are signs of reinfestation, but do not apply more than 0.88 lb ai/A (50 fl oz/A) of TRIPLE CROWN T&O insecticide or 0.50 lb ai imidacloprid/A per year. Repeat application must be limited to no more than once per seven days. Certain cultivars may show sensitivity to the final spray solution. Treat a small number of plants and observe for one week prior to application to the entire planting.			
In Oregon State, do not apply this product, by any application method, to linden, basswood or other Tilia species.			
In New York State, do make a single repeat application of TRIPLE CROWN T&O insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.			

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed, or other pesticides or fertilizers by storage or disposal.

Pesticide Storage

Do not freeze. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

You may contact the Environmental Science U.S., LLC Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Environmental Science U.S., LLC Emergency Response Telephone No. is 1-800-424-9300.

To confine spill: Dike surrounding area or absorb with sand, cat litter, or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Conditions of Sale and Limitation of Warranty and Liability:

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Environmental Science U.S., LLC or Seller. All such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold Environmental Science U.S., LLC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or Environmental Science U.S., LLC, and, to the extent consistent with applicable law, buyer assumes the risk of any such use.

To the extent consistent with applicable law, Environmental Science U.S., LLC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ENVIRONMENTAL SCIENCE U.S., LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ENVIRONMENTAL SCIENCE U.S., LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Owned or Licensed U.S. Patents for this product can be found at <https://patent.envu.com>

Envu and Envu logo are trademarks and TRIPLE CROWN® is a registered trademark owned by Environmental Science U.S., LLC or one of its affiliates.



TRIPLE CROWN[®]

T&O

Insecticide

For use on lawns and landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

envu[™]

Active Ingredients:

	By Wt.
Zeta-Cypermethrin*	2.70%
Bifenthrin**	7.87%
Imidacloprid	13.83%
Other Ingredients***:	75.60%
	100.0%

*Cis/trans isomer ratio: Max 75% ($\pm$) cis and Min. 25% ($\pm$) trans

**Cis isomers 97% minimum; trans isomers 3% maximum.

***Contains Petroleum Distillates

TRIPLE CROWN[®] T&O Insecticide contains 2.25 pounds active ingredients per gallon.

EPA Reg. No. 101563-352

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See other panels for additional precautionary information.

SKU: D00001605
22062313B 250407AV1
Net Contents: 1 Gallon

PRODUCED FOR
Environmental Science U.S., LLC
5000 CentreGreen Way, Suite 400
Cary, NC 27513

PEEL HERE TO OPEN >

