INDAZIFLAM	GROUP	29	HERBICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF: AL, AR, FL, GA, LA, MO, MS, NC, OK, SC, AND TN

Rezilon®

EPA Reg. No. 101563-209

For preemergence control of vaseygrass (Paspalum urvillei) and Dallisgrass (Paspalum dilatatum) from seed in established bermudagrass, bahiagrass, and other warm-season perennial grass pastures and hayfields.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its label.

Control of Vaseygrass and Dallisgrass from Seed

Timing of Application
Rezilon should be applied in late winter, typically through January, for pre-emergence control of Vaseygrass and Dallisgrass *germinating from seed*. Rezilon will not control existing perennial plants or seedlings that have already emerged prior to application. Emerged or established plants should be controlled with appropriate postemergence herbicide. For best results, apply Rezilon in late-winter to allow sufficient soil-incorporating rainfall several weeks prior to weed seed germination. DO NOT apply to water-saturated soil, frozen or snow-covered ground. Avoid applying Rezilon when heavy rainfall is expected as this could move treated soil into areas with desirable vegetation. Do not use flood irrigation to incorporate Rezilon into soil.

Rezilon Use Rates for Established Warm-Season Pastures and Hayfields
Apply Rezilon at 3-5 fl oz/A in late-winter, and no more than 6 fl oz/A in a 12-month period. No single application of Rezilon may exceed 5 fl oz/A. For extended weed control, apply Rezilon at 3 fl oz/A in late-winter followed by another 3 fl oz/A applied soon after the first cutting of hay, or by mid-June in grazed settings. If the rate applied exceeds 3 fl oz/A, hay may not be harvested for a period of 40 days after application. There are no grazing restrictions following an application of Rezilon.

Environmental Science US, LLC.
5000 CentreGreen Way,
Suite 400
Cary, NC 27513

IMPORTANT

Refer to the Rezilon label or other post-emergence herbicide labels for information on adjuvants to be used, rotational restrictions, and additional use directions. The most restrictive provision of any label should apply.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-424-9300.

For PRODUCT USE Information Call 1-800-331-2867).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-800-331-2867

Rezilon is a registered trademark of Environmental Science US LLC.

Environmental Science US, LLC.
5000 CentreGreen Way,
Suite 400
Cary, NC 27513