



A quick and effective way to control house flies

Best uses

Maxforce[®] Fly Spot Bait provides a quick and effective way to control house flies in and on the outside livestock structures, commercial and residential structures, including non-food/non-feed areas of commercial food handling establishments, restaurants, dining areas, storage facilities, schools, hospitals, nursing homes, offices and buildings at marinas. Maxforce Fly Spot Bait may also be applied to the exterior and interior surfaces of dumpsters and other waste containers.

Key strengths

Powerful attractants combined with the active ingredient imidacloprid. This product provides immediate results with a long residual.

Benefits of Maxforce Fly Spot Bait

- Easy application
- Immediate results
- Long residual (up to four weeks when applied indoors)
- Treat less surface area
- Sprayable formulation
- Option to apply as a paint-on treatment in livestock facilities
- Low odour
- Controls pyrethroid-resistant flies

How to apply Maxforce Fly Spot Bait

Dilute Maxforce Fly Spot Bait by mixing 125 g per L of water (1.25%). This is accomplished by adding 2.2 packages per L of water (or 1 package per 450 mL of water).

Paint-on application

Dilution: Add 250 g of Maxforce Fly Spot Bait to 200 mL of water and mix until dissolved.

Use rate: Apply 200 mL of the mixed solution per 100 m² of ground surface area of the structure being treated.

Rate of application for spot treatments

Low to moderate infestations: Apply two litres of finished spray to treat 100 m² of fly resting surfaces.

Moderate to high infestations: Apply four litres of finished spray to treat 100 m² of fly resting surfaces.

Where to apply

Spray diluted Maxforce Fly Spot Bait on fly resting areas such as on walls, around windows, and the exterior and interior surfaces of dumpsters or trash collection bins. Apply product in spots of no greater than 0.2 m² in locations where house flies tend to rest. Spots are not to be adjoining. The combined area of spots is not to exceed 10% of the total surface area of a room.

Directions for paint-on application:

Dilution: Add 250 g of Maxforce Fly Spot Bait to 200 mL of water and mix until dissolved.

Use rate: Apply 200 mL of the mixed solution per 100 m² of ground surface area of the structure being treated.

How to apply: Apply the mixed solution in narrow bands up to 15 cm wide on suitable fly-resting surfaces.

Application equipment: For best results, use a brush or paint roller applicator to apply solution to surfaces. Clean application and mixing equipment with water after application.



Active ingredient
imidacloprid 10%



Group
4A



Formulation
WG



Packaging
50 x 56.7 g packages

Food/feed handling establishments

Applications in food/feed handling establishments are permitted **ONLY** in non-food/non-feed areas. Non-food/non-feed areas include garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets, and storage (after packaging, canning or bottling).

Agriculture Use – Application Instructions

For livestock housing

Where to apply: For use on agricultural livestock production facility structures, such as poultry houses, turkey and chicken houses, cattle, sheep and goat facilities, feedlots, dairy barns, swine confinement buildings, dog kennels, horse stables, animal confinement areas, animal holding areas, animal quarters, animal rearing facilities, livestock barns, hobby and recreational farms, and areas of animal facilities including storage rooms, dumpsters and trash collection areas, where house flies tend to aggregate or be a nuisance.

Why use Maxforce[®] Fly Spot Bait

- Maxforce Fly Spot Bait provides fast and effective results, while eliminating the need to treat large surfaces.
- Maxforce Fly Spot Bait fits well into an overall integrated pest management program for house flies that includes proper sanitation and environmental controls, such as exclusion and proper light management.

Additional application information

- Apply spot treatment applications with low pressure sprayer, do not exceed 20 psi.
- Reapply if house flies continue to be a problem, but not before 7 days.
- Treat as a wettable powder and agitate diluted material frequently throughout the day.
- Applications of this product may leave a visible residue or discolour some surfaces. Before treating, it is advisable to test application in a small inconspicuous area.
- Avoid keeping diluted material in sprayer for more than a couple of days. If storing diluted product in the sprayer, ensure that you properly label and store this material in accordance with provincial and federal pesticide regulations.

Restrictions

- Do not exceed more than 500 g of product per 100 m².
- Treated bait spots must be placed in locations **NOT** accessible to children, pets, livestock or wildlife.
- Do not apply in food areas of food processing or manufacturing facilities or in agriculture facilities.
- Do not apply in classrooms or patient rooms.
- Do not allow people or pets to reenter treated areas until sprays have dried.
- Do not apply to overhead areas or in confined spaces without appropriate respiratory and eye protection.
- Ventilate treated areas after application either by opening windows and doors or through use of air exchange/ventilation systems confirmed to be operational. Use fans where required to aid in the circulation of air.
- Do not allow bait to come in contact with food or water, livestock feed or their water.
- Do not apply to milking equipment at dairy facilities.
- Do not spray directly on animals.
- Where possible, rotate the use of Maxforce Fly Spot Bait or other Group 4 insecticides with different groups that control the same pests.

Darcy Olds

Key Accounts Manager, Professional Pest Management – Canada
darcy.olds@envu.com • 905-464-8399