

11015981

22063326

Fluoxastrobin	GROUP	11	FUNGICIDE
Flutriafol	GROUP	3	FUNGICIDE

TARVECTATM

FUNGICIDE

For the control of foliar, stem and root diseases in turf. For use on turf in golf courses, in lawns and landscape areas around public, industrial, and commercial properties, in athletic fields, and in commercial sod farms.

EPA Reg. No. 279-9706

Active Ingredients:	By Wt.
Fluoxastrobin:	14.84%
Flutriafol:	19.30%
Other Ingredients:	65.86%
TOTAL:	100.0%

Contains 1.40 pounds fluoxastrobin per gallon of (167 g/L) and 1.82 pounds flutriafol per gallon (218 g/L).
Suspension concentrate

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional precautionary information.

M EPA Est. 279-NY-1 CM EPA Est. 86088-PA-001
EPA Establishment Numbers (The letter before each Establishment Number above
corresponds to the first letter in the lot number on the container label.)

Sold by:



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104



00035832559815

FMC, the FMC Logo, and TARVECTATM are
trademarks of FMC Corporation and/or an affiliate.
© 2024 FMC Corporation. All rights reserved.

Net Contents: 2.5 Gallons

GSS-SL-1110 09092024 08-29-2024

FIRST AID

If SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to by a poison control center or doctor.• DO NOT give anything to an unconscious person.
If INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15–20 minutes.• Call a poison control center or doctor for treatment advice.
If IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

For Information Regarding the Use of this Product Call 1-800-321-1FMC(1362).

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals****CAUTION**

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Applicators, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks, and
- Chemical resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber $>$ 14 mils, polyvinyl chloride (PVC) \geq 14 mils or Viton \geq 14 mils.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL or CHEMICAL HAZARDS

DO NOT mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

Environmental Hazards

This pesticide is toxic to mammals, fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. The active ingredient in this product can be persistent for several months or longer. **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, or other sensitive areas that may be exposed to spray drift. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Flutriafol has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In New York State, this product may not be applied within 100 feet of a coastal marsh or stream that drains directly into a coastal marsh. Sale, use, and distribution of this product in Nassau and Suffolk Counties of New York State is prohibited.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water is: coveralls, shoes plus socks, and chemical resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

THE REQUIREMENTS IN THIS BOX APPLY TO USES OF THIS PRODUCT THAT ARE NOT WITHIN THE SCOPE OF THE Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets out of treated area until spray has dried.

PRODUCT INFORMATION

Tarvecta™ fungicide is a broad-spectrum fungicide for the control or suppression of certain diseases in turf. Tarvecta™ fungicide works by interfering with respiration and sterol synthesis in plant-pathogenic fungi and is a potent inhibitor of spore germination and mycelial growth. The active ingredients, fluoxastrobin and flutriafol move rapidly into green tissue via translaminar and acropetal movement. Tarvecta™ fungicide is rainfast 2 hours after application. Disease control will be reduced if rainfall occurs within 2 hours of application. The activity of Tarvecta™ fungicide makes it an excellent choice as a broad spectrum, dual action fungicide for disease management programs.

USE RESTRICTIONS

- **DO NOT** APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.
- **DO NOT** apply this product with aerial application equipment.
- **DO NOT** spray the product into fish pools, ponds, streams, or lakes. **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- **DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- **DO NOT** allow the product to enter any drain during or after application.
- **DO NOT** apply or irrigate to the point of runoff.

RESISTANCE MANAGEMENT

The active ingredients in Tarvecta™ fungicide (fluoxastrobin and flutriafol) belong to the strobilurin (Group 11) and the demethylation inhibitor (Group 3) fungicide classes, respectively. The dual action of Tarvecta™ fungicide provides a built in resistance management strategy that will minimize the resistance in at-risk pathogens.

For resistance management, please note that Tarvecta™ fungicide contains both a Group 11/fluoxastrobin and Group 3/flutriafol fungicide. Any fungal population may contain individuals naturally resistant to Tarvecta™ fungicide and other Group 11 or Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are repeatedly used in the same treated areas. Appropriate resistance-management strategies must be followed. To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Tarvecta™ fungicide or other Group 11 and/or 3 fungicides within a growing season sequence with different groups that control the same pathogens. **DO NOT** make more than two consecutive applications of this product.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact FMC Corporation at 800- 321-1FMC(1382).

APPLICATION GUIDELINES

Broadcast Ground Sprayers

Thorough coverage is necessary to provide good disease control. Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. For ground application equipment 44 gallons/A minimum is required.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate the sprayer before use. Use a pump with the capacity to: (1) maintain a minimum of 35 psi at nozzles, and (2) provide sufficient agitation in the tank to keep the mixture in suspension (this requires recirculation of 10% of the tank volume per minute). Use jet agitators or a liquid sparge tube for vigorous agitation. Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump must be 16-mesh or coarser. **DO NOT** place a screen in the recirculation line. Use 50-mesh screens at the nozzles. Check nozzle manufacturer's directions. For information on spray equipment and calibration, consult sprayer manufacturer's and/or state directions. For specific local directions and spray schedules, consult the current state agricultural experiment station advisements.

Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. **DO NOT** let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Tarvecta™ Fungicide Alone

Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the Tarvecta™ fungicide to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Tarvecta™ fungicide has completely and uniformly dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

Tarvecta™ Fungicide + Tank mix Partners

- Follow the directions of the tank mix partner with respect to adjuvant use.
- **DO NOT** tank mix Tarvecta™ fungicide with any bromoxynil based product or plant injury could occur.

Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, tank mix partners must be added in this order: products packaged in water-soluble packaging (see note in next paragraph), wettable powders, wettable granules, dry flowables, liquid flowables (including Tarvecta™ fungicide), liquids, and emulsifiable concentrates. Always allow each tank mix partner to become fully and uniformly dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

Note: When using Tarvecta™ fungicide in tank mixtures, all products in water-soluble packaging must be added to the tank before any other tank mix partner. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If using Tarvecta™ fungicide in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations, which appear on the tank mix product label. No label dosage rate may be exceeded, and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product that prohibits such mixing. Tank mixtures or application of other products referenced on this label are permitted only in those states in which the referenced products are registered.

Compatibility

Tarvecta™ fungicide is physically compatible with most insecticide, fungicide, herbicide and foliar nutrient products. However, the physical compatibility of Tarvecta™ fungicide with tank mix partners must be tested before use. To determine the physical compatibility of Tarvecta™ fungicide with other products, use a jar test, as described in the next paragraph.

Jar Test Procedure: Using a quart jar, add the proportionate amounts of the products to ½ qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, add the remaining ½ qt of water, shake, and let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Mandatory Spray Drift Management

Ground Boom Applications:

- **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.
- Select a nozzle and pressure combination that delivers medium to coarser droplet size (ASAE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to use a medium to coarser droplet size (ASAE S572) for all applications.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Boomless Ground Applications

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

- Take precautions to minimize spray drift.

Controlling Droplet Size – Ground Boom

Volume: Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow volume.

Pressure: Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.

Spray Nozzle: Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom: For ground equipment, the boom must remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce the effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed.
Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Sensitive Areas

This pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non- target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

APPLICATION TO TURF

Tarvecta™ fungicide provides control or suppression of several important diseases listed for turf. Use Tarvecta™ fungicide in conjunction with cultural practices that promote healthy, vigorous turfgrass. These practices include nutrient management, thatch management, water management and judicious use of other fungicides and cultural practices.

TURF USE DIRECTIONS

Uniform application is necessary to provide good disease control. For soil-borne diseases, use sufficient water to move the active ingredient into the crown and upper root zone, or irrigate after application with 1/8" to 1/4" water.

Use Rate and Retreatment Interval Ranges: Under light to moderate disease pressure, use lower specified use rates and/or longer specified retreatment intervals. Under conditions of high disease pressure or when prolonged favorable disease conditions exist, use shorter specified retreatment intervals and/or higher specified use rates.

Turfgrass Use Rate Conversion				
Fl oz product/ 1000 sq ft	lb ai flutriafol/ A	lb Fluoxastrobin/ A	Total AI/ A	Fl oz product/ A
0.35	0.21	0.16	0.38	15
0.70	0.43	0.33	0.76	30

DIRECTIONS FOR APPLICATION TO TURF

Disease Control		Product Rate		Application Interval (Days)
		fl oz/ 1000 sq ft	fl oz/ A	
Apply preventatively when conditions are favorable for disease development	Anthrachnose Basal and Foliar (<i>Colletotrichum cereale</i>) ¹	0.35 - 0.70	15 - 30	14-21
	Brown Patch (<i>Rhizoctonia solani</i>) ²			14-28
	Brown Ring Patch (<i>Waitea circinata</i> var. <i>circinata</i>) or (Waitea patch) ² ¥			14-28
	Dollar Spot (<i>Clavireedia</i> sp.)			14-21
	Gray Leaf Spot (<i>Pyricularia grisea</i>)			14-21
	Leaf Spot (<i>Bipolaris sorokiniana</i>)			14-21
	Melting Out (<i>Drechslera poae</i>)			14-21
	Mini Ring (<i>Rhizoctonia zeae</i>)			14-21
	Necrotic Ring Spot (<i>Leptosphaeria korrea</i>) ¥			21-28
	Pythium Root Dysfunction (<i>Pythium volutum</i>) ³			28
	¹ Under high disease pressure, use the higher rate. ² For longer residual use higher rate. ³ Apply in fall and spring when mean daily soil temperatures are between 50 - 75° F at a 2” depth.			
Apply preventatively in fall	Large Patch (<i>Rhizoctonia solani</i>) ⁴	0.35 - 0.7	15 - 30	28
	Spring Dead Spot (<i>Ophiosphaerella korrea</i> , <i>O.narmari</i> , <i>O. herpotricha</i>) ⁵			
⁴ Apply when soil temperatures at a 2” depth are below 70° F. One repeat application may be made 28 days after initial treatment in fall. Spring applications must be made when soil temperatures at a 2” depth are consistently between 50 – 55° F. ⁵ Apply initial application when soil temperatures at a 2” depth are between 70 – 75° F. Incorporate treatment with post-application irrigation.				

(table continued on next page)

Disease Control		Product Rate		Application Interval (Days)
		fl oz/ 1000 sq ft	fl oz/ A	
Apply preventatively in fall and/or spring	Take-All Patch (<i>Gaeumannomyces graminis f. sp. avenae</i>) ⁶	0.35-0.7	15-30	28
	Take-All Root Rot (Multiple species) ⁷			21-28
	Yellow Patch (<i>Rhizoctonia cerealis</i>) ⁸			28
	⁶ Apply preventatively in fall when soil temperatures are between 55 – 60° F. Repeat application must be made in spring when soil temperatures are favorable for pathogen activity. ⁷ Apply preventatively in fall when soil temperatures are between 75 – 80° F. Repeat applications are advised in spring once soil temperatures are between 55 – 60° F. ⁸ Make one or two applications in fall or when conditions are favorable for disease development. Applications may be made in the spring if disease appears.			
Apply in Spring	Fairy Ring (Multiple species) ⁹	0.35-0.7	15-30	28
	Summer Patch (<i>Magnaporthe poae</i>) ¹⁰			21-28
	⁹ Applications must begin when soil temperatures at a 2" depth are 55 - 60°F, or as soon as visual disease symptoms are observed. Sufficient spray volume (at least 2 gal/1000 sq ft) is needed to move fungicide down through thatch layer where pathogen is active. Post-application irrigation or addition of a wetting agent is advised to incorporate fungicide. ¹⁰ Apply preventatively in spring when soil temperature reaches 65° F at a depth of 2".			
Apply preventative application prior to permanent snow cover	Gray Snow Mold/Typhula Blight (<i>Typhula incarnata</i>) ¹¹	0.35-0.7	15-30	28
	Pink Snow Mold (<i>Microdochium nivale</i>) ¹¹			
	¹¹ If two applications are needed, make the first application at the beginning of turfgrass dormancy and another one just prior to permanent snow cover.			

(table continued on next page)

Restrictions:

- No single application may exceed 35 fl oz of product/A (0.50 lb flutriafol/A and 0.38 lb fluoxastrobin/A).
- **DO NOT** apply more than 3 applications per year.
- The minimum reapplication interval is 14 days. Refer to the "application interval" directions for disease-specific details.
- **DO NOT** apply more than 105 fl oz of product (0.82 gals of product)/ A per year.
- **DO NOT** apply more than 1.5 lb ai of flutriafol/A per year.
- **DO NOT** apply more than 2.13 lb ai of fluoxastrobin/A per year.
- **DO NOT** use on home lawns (i.e., residential turf) and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- **DO NOT** use clippings for animal feed.

¥ Not registered for use in California.

For use in the establishment of turfgrass from seed or in overseeding of dormant turfgrass:

Tarvecta™ fungicide may be used for control of certain turfgrass diseases associated with turfgrass establishment from seed. Tarvecta™ fungicide may also be used during overseeding of dormant turfgrass. Tarvecta™ fungicide may be safely applied before or after seeding or at seedling germination and emergence to ryegrass, bentgrass, bluegrass, fescue, and other turfgrasses. Optimum application timing for control of seeding diseases is just prior to, during or just after seeding.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed. Store in a cool, dry, secure place. In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills. For assistance call: 800-424-9300.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Rigid, Nonrefillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons).

Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.



11015981

22063326

Fluoxastrobin	GROUP	11	FUNGICIDE
Flutriafol	GROUP	3	FUNGICIDE

TARVECTATM

FUNGICIDE

For the control of foliar, stem and root diseases in turf. For use on turf in golf courses, in lawns and landscape areas around public, industrial, and commercial properties, in athletic fields, and in commercial sod farms.

EPA Reg. No. 279-9706

Active Ingredients:

	By Wt.
Fluoxastrobin:	14.84%
Flutriafol:	19.30%
Other Ingredients:	65.86%
TOTAL:	100.0%

Contains 1.40 pounds fluoxastrobin per gallon of (167 g/L) and 1.82 pounds flutriafol per gallon (218 g/L).
Suspension concentrate

KEEP OUT OF REACH OF CHILDREN
CAUTION

See inside booklet for additional precautionary information.

M EPA Est. 279-NY-1

CM EPA Est. 86088-PA-001

EPA Establishment Numbers (The letter before each
Establishment Number above corresponds to the first letter
in the lot number on the container label.)

Sold by:



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104



00035832559815

Net Contents: 2.5 Gallons

GSS-SL-1110 09092024 08-29-2024

OPEN HERE ▲

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.








THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.

Dieline does not print.



Cimarron Label
inovar

4201 North Westport Ave. • Sioux Falls, SD 57107
Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
10/18/24	241266	FMC
LABEL SIZE	BOOKLET SIZE	
6.75" x 6.75"	6.75" x 6"	
LABEL COLORS	BOOKLET OUTSIDE COLORS	BOOKLET INSIDE COLORS
<div> BLK</div> <div> 032</div> <div> 369</div>	<div> BLK</div> <div> 032</div> <div> 369</div>	<div> BLK</div>
PATTERN VARNISH: <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO		

Form: CS 006B - 3/29/2017

☐ **ARTWORK IS APPROVED**

☐ **REVISED PROOF NEEDED**

**WE CANNOT PROCESS
THIS ORDER WITHOUT AN
AUTHORIZED SIGNATURE**

Signed _____ **Date** _____