



Environmental Science US, LLC  
5000 CentreGreen Way, Suite 400  
Cary, NC 27513

**Resilia™**

**For control of Mini Ring in turfgrass.**

**EPA Reg. No. 101563-223**

## ***FIFRA 2(ee) Recommendation***

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF ALABAMA, ARIZONA, ARKANSAS, FLORIDA, GEORGIA, HAWAII, KANSAS, KENTUCKY, LOUISIANA, MARYLAND, MISSISSIPPI, MISSOURI, NEBRASKA, NEVADA, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS, VIRGINIA**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**DIRECTIONS FOR USE:** It is a violation of federal law to use this product in a manner inconsistent with its label. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**TURF USE DIRECTIONS:** RESILIA controls a range of turf root diseases. Refer to the table below for target pest, rate structure, and specific application instructions. Apply RESILIA preventively when conditions are favorable for target pest activity. Reapply as needed, using shorter intervals when target pest pressure is high but Do Not exceed maximum specified rate. Adequately water-in applications to the depth of the root zone or target pest. See 'Application Directions' for further application instructions on equipment, nozzles, spray volumes and mixing instructions.

Target Pest	Application Rate (fl oz Product /1000 ft <sup>2</sup> ) <sup>1</sup>	Interval between Applications (days)	Use Information
Mini Ring (caused by <i>Rhizoctonia zeae</i> )	4	14-28	Apply Resilia preventively when conditions are favorable for target pest activity. Reapply as needed, using shorter intervals when target pest pressure is high but do not exceed maximum specified rate. Adequately water-in applications to the depth of the root zone or target pest.

<sup>1</sup>4 fl oz product/1,000 ft<sup>2</sup> equates to 0.11 lbs a.i./A of fluopyram, 0.2 lbs a.i./A of prothioconazole, and 4.1 lbs a.i./A of propamocarb hydrochloride.

### **RATE RESTRICTIONS**

- 1) **Maximum Single Rate:** Do not exceed 4 fl oz product/1000 ft<sup>2</sup> per application, equivalent to 0.11 lbs a.i./A of fluopyram, 0.2 lbs a.i./A of prothioconazole, and 4.1 lbs a.i./A of propamocarb hydrochloride.
- 2) **Yearly Maximum Rate:** Do not exceed 16 fl oz product/1000 ft<sup>2</sup> per year, equivalent to 0.44 lbs a.i./A/year of fluopyram, 0.8 lbs a.i./A/year of prothioconazole, and 16.4 lbs a.i./A/year of propamocarb hydrochloride.
- 3) Do not exceed 4 applications per year using the maximum single application rate.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-331-2867.

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-800-331-2867.*