



FLUOPYRAM	GROUP	7	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF AK, AL, AR, AZ, CA, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NY, NV, OH, OK, OR, PA, PR, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

BROADFORM®

EPA Reg. No. 101563-158

For Control of Beech Leaf Disease on Beech Trees

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

Plant	Target Disease	Application Rate	Interval between applications (days)
Beech Tree	Beech Leaf Disease (BLD)	4 - 8 fluid ounces/100 gallons	7 - 21 until the threat of disease is over
	<ul style="list-style-type: none">Do not apply more than 27.3 fl. oz. of BROADFORM per acre per year.Applications are to be made through hydraulic sprayers operating between 150-350 psi.Use of a spreader sticker is recommended to improve efficacy for BLD.		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-424-9300.

For PRODUCT USE Information Call 1-800-331-2867.

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-800-331-2867.

**Environmental Science US, LLC.
5000 CentreGreen Way, Suite 400
Cary, NC 27513**