


Product Information Sheet



- Active Ingredient: Fosetyl-Al 80%
- FRAC Group: P07
- EPA Registration Number: 432-890
- Signal Word: Caution
- Formulation: Wettable Dispersible Granule
- REI: 12 hours



Aliette® provides outstanding systemic control of Phytophthora and Downy Mildew diseases.

Best Uses

- Effective on economically-important water mold diseases
- Controls *Phytophthora*, Downy Mildew and Pythium
- Control of Fire Blight and suppression of Bacterial Blight caused by certain pathovars of *Xanthomonas campestris*
- Rotation and resistance management in settings where other products have been overused


Attributes

- True systemic movement in the plant xylem and phloem, protecting existing and new plant tissue
- Up to 30 days of insect control
- Stimulates plant's disease defense mechanisms
- No documented cases of resistance
- Tank-mixes readily with other fungicides for a wider spectrum of control
- Do not apply within 14 days of copper-based treatments

Application Rates

- For foliar applications, apply at 2.5-5 lb./100 gal. of water
- For drench applications, apply at 6.4-12.8 oz./100 gal. of water
- Aliette WDG is available in 5-lb. containers

Downy Mildew Control on Stock



Dr. Ann Chase, Chase Horticultural Research, 2002

Ratings were made 37 Days After Application (DAA)

Phytophthora Root and Crown Rot on Calibrachoa



Fungicidal Control of Phytophthora Root and Crown Rot on Calibrachoa
Margery Daughtrey, Cornell, 2007

Ratings were made 33 Days After Application (DAA)

* Root Rot rated on a 1-4 scale, with 1=best roots, 2=some Root Rot, 3=Root Rot with some white roots still visible and 4=complete Root Rot.